

University of Groningen

Stars, neutral hydrogen and ionised gas in early-type galaxies

Serra, Paolo

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2008

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Serra, P. (2008). *Stars, neutral hydrogen and ionised gas in early-type galaxies*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



rijksuniversiteit
 groningen

Stars, neutral hydrogen and ionised gas in early-type galaxies

Proefschrift

ter verkrijging van het doctoraat in de
 Wiskunde en Natuurwetenschappen
 aan de Rijksuniversiteit Groningen
 op gezag van de
 Rector Magnificus, dr. F. Zwarts,
 in het openbaar te verdedigen op
 vrijdag 11 april 2008
 om 16.15 uur

door

Paolo Serra

geboren op 18 maart 1980
 te Cagliari, Italië

Promotor:
Copromotores:

Prof. dr. S. C. Trager
Dr. T. A. Oosterloo
Dr. R. Morganti

Beoordelingscommissie:

Prof. dr. F. Combes
Prof. dr. J. H. van Gorkom
Prof. dr. R. F. Peletier

Cover – H I gas around NGC 4278 with column density above $1 \times 10^{19} \text{ cm}^{-2}$. Observations were obtained by Raffaella Morganti with the Westerbork Synthesis Radio Telescope.

Contents

| | | |
|----------|---------------------------------------------------------------------------------|-----------|
| 1 | Introduction | 9 |
| 1.1 | Observational facts about nearby early-type galaxies | 10 |
| 1.1.1 | Scaling relations | 10 |
| 1.1.2 | Stellar populations | 11 |
| 1.1.3 | Structure | 12 |
| 1.1.4 | Gas | 13 |
| 1.2 | Formation and evolution of early-type galaxies | 15 |
| 1.2.1 | Monolithic collapse and galaxy merging | 15 |
| 1.2.2 | Evidence of merging activity | 16 |
| 1.2.3 | The role of gas | 17 |
| 1.2.4 | Accretion from the intergalactic medium | 18 |
| 1.3 | This thesis | 19 |
| 1.3.1 | Relation between gas and stellar properties | 19 |
| 1.3.2 | H I and goals of this work | 20 |
| 1.3.3 | Outline of the thesis | 21 |
| 2 | An H I-selected early-type galaxy sample | 23 |
| 2.1 | Sample selection | 23 |
| 2.1.1 | HIPASS and the southern sample | 23 |
| 2.1.2 | Northern sample | 24 |
| 2.2 | Sample properties | 30 |
| 2.3 | H I-rich galaxies | 33 |
| 2.4 | Summary | 48 |
| 3 | The case of IC 4200: stellar, H I and ionised gas content and kinematics | 49 |
| 3.1 | Basic Properties of IC 4200 | 50 |
| 3.2 | Radio interferometry | 52 |
| 3.2.1 | Observations | 52 |
| 3.2.2 | H I | 52 |
| 3.2.3 | Continuum | 53 |
| 3.3 | Optical Spectroscopy | 53 |
| 3.3.1 | Observations | 53 |

| | | |
|--------------|------------------------------------------------------------------------------------------------------------------------------|------------|
| 3.3.2 | Disentangling stars from ionised gas | 56 |
| 3.3.3 | Absorption-line indices | 57 |
| 3.4 | Optical Imaging | 58 |
| 3.5 | Results | 59 |
| 3.5.1 | H I | 59 |
| 3.5.2 | Radio continuum | 63 |
| 3.5.3 | Stellar and ionised gas kinematics | 64 |
| 3.5.4 | Gas ionisation | 65 |
| 3.5.5 | Stellar populations | 66 |
| 3.5.6 | Optical Imaging | 68 |
| 3.6 | Discussion | 70 |
| 3.6.1 | Origin of the H I disk: merger or IGM accretion? | 70 |
| 3.6.2 | Clues from the stellar properties | 72 |
| 3.6.3 | The fate of IC 4200 | 75 |
| 3.7 | Conclusions | 75 |
| Appendix 3.A | Calibration of the line-strength indices to the Lick/IDS system | 77 |
| Appendix 3.B | Velocity dispersion corrections of the line-strength indices | 78 |
| Appendix 3.C | Tables | 78 |
| 4 | Stellar populations, neutral hydrogen and ionised gas in early-type galaxies | 83 |
| 4.1 | Observations, data reduction and analysis | 83 |
| 4.1.1 | Separating stellar spectrum from ionised-gas emission lines | 84 |
| 4.1.2 | Lick/IDS line-strength indices and derivation of t_{SSP} , $[Z/H]_{\text{SSP}}$ and $[E/Fe]_{\text{SSP}}$ | 85 |
| 4.2 | Results | 87 |
| 4.2.1 | Ionised gas: properties and connection to the H I content | 88 |
| 4.2.2 | Stellar populations | 89 |
| 4.2.3 | Stellar populations and H I | 92 |
| 4.2.4 | H β correction and the case of IC 4200 | 92 |
| 4.3 | Discussion | 95 |
| 4.3.1 | Ionised gas | 95 |
| 4.3.2 | Stellar populations | 95 |
| 4.3.3 | Mergers of galaxies | 96 |
| 4.3.4 | Structural properties and IGM accretion | 98 |
| 4.4 | Conclusions | 99 |
| Appendix 4.A | Calibration of the line-strength indices to the Lick/IDS system | 108 |
| 5 | Neutral hydrogen and the kinematics of stars and ionised gas in early-type galaxies | 113 |
| 5.1 | Three H I-rich southern galaxies | 114 |
| 5.1.1 | ATCA observations and data reduction | 114 |
| 5.1.2 | ESO 092-21 | 115 |
| 5.1.3 | ESO 140-31 | 117 |
| 5.1.4 | ESO 381-47 | 119 |
| 5.2 | Optical spectroscopy: stellar and ionised-gas kinematics | 122 |
| 5.2.1 | Data analysis | 122 |

| | | |
|----------|-------------------------------------------------------------------------------------------------------------------------------------|------------|
| 5.2.2 | Error analysis | 122 |
| 5.2.3 | Results | 123 |
| 5.2.4 | Comparison to previous results | 145 |
| 5.3 | Comparison to H I properties | 146 |
| 5.3.1 | Stellar kinematics, H I and the v/σ vs. ϵ diagram | 146 |
| 5.3.2 | Ionised-gas and H I kinematics | 150 |
| 5.4 | Stellar kinematics and populations | 151 |
| 5.5 | Summary | 153 |
| | Appendix 5.A Kinematics of individual galaxies | 154 |
| 6 | On the interpretation of age and chemical composition of composite stellar populations determined with line-strength indices | 165 |
| 6.1 | Introduction | 166 |
| 6.2 | Models of composite stellar populations | 167 |
| 6.3 | Results | 169 |
| 6.4 | Conclusions | 173 |
| 7 | Summary and future prospects | 177 |
| 7.1 | Results of this thesis | 178 |
| 7.1.1 | H I properties of ETGs | 178 |
| 7.1.2 | Individual galaxies: the case of IC 4200 | 178 |
| 7.1.3 | H I and stellar populations | 179 |
| 7.1.4 | H I and stellar kinematics | 180 |
| 7.1.5 | H I and ionised gas | 181 |
| 7.2 | Limitations of this thesis | 182 |
| 7.3 | Future work: Atlas ^{3D} and a WSRT H I survey | 183 |
| | Bibliography | 185 |
| | Nederlandse samenvatting | 197 |
| | Acknowledgments | 205 |

